



Management Plan

For Antarctic Specially Protected Area (ASPA) No. 161 TERRA NOVA BAY, ROSS SEA

Introduction

The ASPA of Terra Nova Bay is a coastal marine area encompassing 29.4 km² between Adélie Cove and Tethys Bay, Terra Nova Bay, immediately to the south of the Italian Mario Zucchelli Station (MZS). Terra Nova Bay was originally designated as Antarctic Specially Protected Area through Measure 2 (2003) after a proposal of Italy. CCAMLR considered and approved its designation during CCAMLR XXVI, Hobart 2007. The Management Plan has been revised in 2008, through measure 14 (2008) and in 2013 through measure 15 (2013).

The primary reason for the designation of Terra Nova Bay as an Antarctic Specially Protected Area (ASPA) is its particular interest for ongoing and future research. Long term studies conducted in the last 30 years by Italian scientists have revealed a complex array of species assemblages, characterized by unique symbiotic interactions. In this Area, several VME species are also present, above all the Antarctic scallop *Adamussium colbecki* and pterobranchs, and new species continue to be described.

The high ecological and scientific values derived from the diverse range of species and assemblages, together with the vulnerability of the Area to disturbance by scientific oversampling, alien introductions, and direct human impacts arising from increasing activities at the nearby permanent scientific stations (also considering the construction of the new gravel runway at Boulder Clay - Final CEE, 2017) are such that the Area requires long-term special protection.

No Domain nor ACBR number is proposed as the Environmental Domain Analysis for Antarctica (Resolution 3, 2008) and Antarctic Conservation Biogeographic Regions (Resolution 6, 2012) classifications are based on terrestrial criteria.

1. Description values to be protected

This coastal marine area is an important area for well-established and long-term scientific investigations that allowed, up to now, to collect an extensive amount of scientific data. The site typically remains ice-free in summer, which is rare for coastal areas in the Ross Sea region, making it an ideal and accessible site for research into the near-shore benthic communities of the region. Extensive marine ecological research has been carried out at Terra Nova Bay since 1986/87, contributing substantially to our understanding of the marine communities in this area, and of the effect of katabatic winds on the physical, chemical and biological processes occurring in the water column (Povero et al., 2001).

High diversity at both species and community levels make this Area of high ecological and scientific value. Studies have revealed a complex array of species assemblages, often co-existing in mosaics (Sarà et al., 1992; 2002; Gambi et al., 1997; Cantone et al., 2000; Ghiglione et al., 2013) and characterized by unique symbiotic interactions (Schiaparelli et al., 2011; 2015; Regoli et al., 2004). There exist assemblages with high species richness and complex functioning, such as the sponge and anthozoan communities, alongside loosely structured, low diversity assemblages. In this area several VME species also occur, above all the Antarctic scallop *Adamussium colbecki* (Schiaparelli and Linse, 2006) and pterobranchs (Schiaparelli et al., 2004), and new species continue to be described (Schiaparelli and Jirkov, 2016). A population of Adélie penguins (*Pygoscelis adeliae*) is present nearby the Area.

The collected scientific data over the years, allowed the site to serve as reference for the determination of impacts arising from human activities (Berkman and Nigro, 1992; Focardi et al., 1993; Minganti et al., 1995; Bruni et al., 1997; Nonnis Marzano et al., 2000, Lo Giudice et al., 2013).



2. Aims and objectives

- avoid degradation of, or substantial risk to, the values of the Area by preventing unnecessary human disturbance to the Area;
- allow scientific research on the ecosystem, in particular on the marine species assemblages and long-term monitoring, while ensuring protection from oversampling or other possible human impacts;
- allow other scientific research and support activities provided they are for compelling reasons which cannot be served elsewhere;
- minimise the possibility of introduction of alien plants, animals and microbes to the Area;
- allow visits for management purposes in support of the aims of the Management Plan.

3. Management activities

The following management activities are to be undertaken to protect the values of the Area:

- A map showing the location of the Area (stating the special restrictions that apply) shall be displayed, and a copy of this Management Plan shall be kept available, at all the scientific stations located within 50 km of the Area. Information illustrating the location and boundaries with clear statements of entry restrictions is displayed on Posters at MZS;
- Buoys, or other markers or structures established for scientific or management purposes shall be secured and maintained in good condition, and removed when no longer necessary;
- Any abandoned equipment or material shall be removed to the maximum extent possible, provided that doing so does not adversely impact on the environment and the values of the Area;
- Visits shall be made as necessary to assess whether the Area continues to serve the purposes for which it was designated and whether management and maintenance measures are adequate.
- National Antarctic Programs are encouraged to consult together to prevent oversampling within the Area.

4. Period of designation

Designated for an indefinite period.

5. Maps and photographs

Map 1: Terra Nova Bay, Antarctic Specially Protected Area No. 161, bathymetric map.

Map specifications: Projection: UTM Zone 58S; Spheroid: WGS84. Bathymetric contour interval 50 m. Land contours and coast derived from 1:50,000 Northern Foothills Satellite Image Map (Frezzotti *et al.* 2001). Bathymetry within ASPA derived from high resolution sidescan sonar data surveyed by Kvitek, 2002. Bathymetry outside of ASPA supplied by Italian Hydrographic Office 2000. Marine data collected under Terra Nova Bay marine protected area Project (PNRA 1999–2001). Inset 1: The location of Terra Nova Bay in Antarctica. Inset 2: Terra Nova Bay location map, showing the region covered by Map 1, stations, and sites of nearby protected areas.

6. Description of the Area

6(i) Geographical co-ordinates, boundary markers and natural features

General description, borders and coordinates

The Area is situated in Terra Nova Bay, between the Campbell Glacier Tongue and Drygalski Ice Tongue, Victoria Land (Map 1). It is confined to a narrow strip of coastal waters to the south of MZS (Italy), extending approximately 9.4 km in length and generally within 1.5 – 7 km of the shore, comprising an area of 29.4 km² (Map 1). No marine resource harvesting has been, is currently, or is planned to be conducted within the Area, nor in the immediate surrounding vicinity.

The western boundary of the Area is defined as the mean high water mark along the coastline extending between 74°42'57"S in the north (2.3 km south of MZS) and 74°48'00"S in the south (the southern shore of Adélie Cove), and includes the intertidal zone (Map 1). The northern boundary of the Area is defined as the 74°42'57"S line of latitude, extending from the coast 1.55 km eastward to the 164°10'00"E line of longitude. The boundary position may be recognised near the shore by the presence of a large and distinctive offshore rock in the northernmost cove on the coast south of MZS, which is an unique feature on this stretch of coast. The southern boundary is defined as the 74°48'00"S line of latitude, extending from the coast 3.63 km eastward to the 164°10'00"E line of longitude. The boundary position may be recognized visually as being at the southern shore of the mouth of Adélie Cove, immediately south of a distinctive rocky outcrop at the base of the coastal cliffs. The eastern boundary of the Area is defined as the 164°10'00"E line of longitude extending between 74°42'57"S in the north and 74°48'00"S in the south.



Geology

The coastline of Terra Nova Bay is characterised predominantly by rocky cliffs, with large boulders forming occasional 'beaches' (Simeoni *et al.*, 1989). In the sheltered areas, the soft bottom begins at a depth of 20–30 m. The tidal range is 1.5–2 m and pack ice approximately 2–2.5 m thick covers the sea surface for 9–10 months of the year (Stocchino and Lusetti, 1988; 1990). Data available for the summer period suggest that ocean currents in the Area are likely to be slow and to flow generally in a north-south direction. Along the coastline of the Area there are two main coves; the larger Adélie Cove in the south and a smaller cove around 3 km to its north. The sea floor substrate of the smaller consists of pebbles of various sizes, while Adélie Cove is characterised by fine-grained, muddy sediments. The seafloor within the Area is primarily granitic rock, with softer substrates composed of coarse sands or gravels.

Invertebrates (0–40 m)

In the supralittoral zone, only cyanobacteria and diatoms colonise the hard substrates, while the intertidal zone (1.5–2.0 m wide) has, in the most sheltered areas, a high coverage of the green alga *Urospora penicilliformis* and *Prasiola crispa* (Cormaci *et al.*, 1992b). Below the tidal zone, down to 2–3 m depth, the community is very poor, due to the persistent presence and scouring action of pack ice, and is mainly composed of epilithic diatoms and the crustacean amphipod *Paramoera walkeri*. Immediately deeper, rocks can be fully colonised by the red alga *Iridaea cordata* (Cormaci *et al.*, 1996), frequently found with *Plocamium cartilagineum*, to a depth of 12 m (Gambi *et al.*, 1994; 2000a). At this level, large sessile animals such as *Alcyonium antarcticum* and *Urticinopsis antarctica* can be occasionally observed, while frequent are the asteroid *Odontaster validus* and the echinoid *Sterechinus neumayeri*. *Phyllophora antarctica* is another red alga forming expanded mats from 12 to 25 m depth, often fully colonised by sessile organisms, mainly hydroids (Cerrano *et al.*, 2000c, Puce *et al.*, 2002), serpulids and bryozoans (*Celleporella antarctica* and *Harpecia spinosissima*). The upper algal belts represent shelter and a food source for diversified and abundant communities of mobile fauna. Numerous invertebrates, such as the polychaete *Harmothoe brevipalpa*, the mollusc *Laevilitorina antarctica*, the crustacean amphipod *Paramoera walkeri* and the isopod *Nototanaïs dimorphus* feed on these algal species and can be very abundant. On rocky bottoms in deeper layers, the algal colonisation is replaced by a calcareous crustose coralline alga (*Clathromorphum lemoineanum*) on which sea-urchins feed.

The soft bottoms from 20–40 m depth are coarse sands and gravels, where the community is characterised by the mollusc bivalve *Laternula elliptica* and the polychaete *Aglaophamus ornatus* (Nephtidae). The bivalve *Yoldia eightsi* is abundant in fine-sand sediments especially in the Adélie Cove.

Invertebrates (30–70 m)

Between 30–70 m, the substrate becomes finer and is completely colonised by the bivalve *Adamussium colbecki*, the shells of which are colonised by a micro-community comprising mainly forams, bryozoans (*Aimulosia antarctica*, *Arachnopusia decipiens*, *Ellisina antarctica*, *Micropora brevissima*) and the spirorbid *Paralaeospira levinsenii* (Albertelli *et al.*, 1998; Ansell *et al.*, 1998; Chiantore *et al.*, 1998; 2000; 2001; 2002; Vacchi *et al.*, 2000a; Cerrano *et al.*, 2001a; 2001b). In this region, large predators such as the gastropod *Neobuccinum eatoni* and the nemertean *Parborlasia corrugatus* are frequent. The echinoid *Sterechinus neumayeri* and the starfish *Odontaster validus* are still very frequent at all depths on both hard and mobile substrates (Chiantore *et al.*, 2002; Cerrano *et al.*, 2000b). Several unique biotic associations have been described at these depths, e.g. between sponges and other invertebrates (Schiaparelli *et al.*, 2000; 2003; 2007; 2010; 2011; 2015). Sponge also represent a key taxon, which has been widely investigated in terms of symbionts (Regoli *et al.*, 2004) and associated microbes (Lo Giudice *et al.*, 2019). In recent years also new species for science have been described as the parasitic amphipod *Lepidepecrella debroyeri* (Schiaparelli *et al.*, 2015) and the ampharetid polychaete *Amphicteis teresae* (Schiaparelli and Jirkov, 2016). Other data have been produced about VME species, such as *Cephalodiscus densus* (Schiaparelli *et al.*, 2004) and *Adamussium colbecki* (Schiaparelli and Linse, 2006). About the latter species, new analyses of data collected in 2006–2007, thank to the presence of a mooring within the ASPA boundaries (Mooring "L" under the Italian mooring code system), showed that this species recruits during summer months in coincidence with an increase of the seawater temperature and a seasonal shift in the water currents and intensity (Schiaparelli and Aliani, 2019).

Invertebrates (70–130 m)

Below 70–75 m down to 120–130 m depth, heterogeneous substrates allow hard- and soft-bottom communities to coexist. On the sparse rocky outcrops the encrusting algae disappear and the benthic communities are dominated by the sessile zoobenthos. This diversified filter feeding assemblage is mainly characterised by sponges and anthozoans, while in soft sediments detritus-feeder polychaetes and bivalves dominate. Among sponges, which can reach very high biomass values, *Axociella nidificata*, *Calyx arcuarius*, *Gellius rudis*, *Phorbas glaberrima*, *Tedania charcoti*, are very abundant (Sarà *et al.*, 1992; 2002; Gaino *et al.*, 1994; Cattaneo-Vietti *et al.*, 1996; 2000c; Bavestrello *et al.*, 2000; Cerrano *et al.*, 2000a). Numerous invertebrates constitute an important component of this assemblage which develops down to 120–140 m depth. These include crustacean peracarids, pycnogonids, mollusc opisthobranchs (*Austrodoris kerguelenensis*, *Tritoniella belli*) (Cattaneo-Vietti, 1991; Gavagnin *et al.*, 1995) and bivalves, ophiuroids and holothuroids, bryozoans, and a variety of endobionts. The conspicuous sponge spicule mats found at these depths underline the important role of sponges in this area, besides the one played by diatoms, in determining the sediment texture and silica content. A peculiar community, dominated by polychaetes and by the bivalve *Limatula hodgsoni*, can be associated with these mats.

Invertebrates (below 130 m)

Below 130 m the hard substrates become very sparse and are mainly colonised by the polychaete *Serpula narconensis* (Schiaparelli *et al.*, 2000) and several bryozoans (*Arachnopusia decipiens*, *Ellisina antarctica*, *Flustra angusta*, *F. vulgaris* and *Isoschizoporella similis*). The dominant muddy bottoms are instead characterised by tubicolous polychaetes (Gambi *et al.*, 2000b), mainly *Spiophanes*. Much deeper, at about 150–200 m depth, brachiopods and various species of bivalves characterise the environment on small gravels as well as on the soft bottom (Cattaneo-Vietti *et al.*, 2000b). The great heterogeneity of these substrates contributes to the creation of communities with considerable species richness, diversity and biomass.



Bird, fish and mammals

An Adélie penguin (*Pygoscelis adeliae*) colony is situated nearby the Area at Adélie Cove, with a 2013 population of 13,408 breeding pairs (Humphries *et al.*, 2017) (Map 1). About 30 Skua (*Stercorarius maccormicki*) pairs breed close to the penguins (Final CEE – 2017).

The faunal assemblage of the Area includes notothenioid fishes, represented especially by species of the *Trematomus* group, including *T. bernacchi*, *T. pennelli*, *T. hansonii* and *T. loennbergii*. These exert an important role in benthic food webs as consumers of many invertebrate species, mainly crustaceans and polychaetes (Vacchi *et al.*, 1991; 1992; 1994a; 1994b; 1995; 1997; 2000b; La Mesa *et al.*, 1996; 1997; 2000; Guglielmo *et al.*, 1998). The platelet ice occurring at Terra Nova Bay in early spring has been shown to house an important nursery for the Antarctic silverfish, *Pleuragramma antarcticum*, a key organism in the ecology of Antarctic food webs (La Mesa *et al.*, 2004; Vacchi *et al.*, 2004). The platelet ice environment has strong prooxidant characteristics at the beginning of austral spring, and the marked responsiveness of antioxidant defences represents a fundamental strategy for *P. antarcticum* (Regoli *et al.*, 2005b).

An aerial survey on cetacean species, conducted in the coastal area surrounding the Italian Station Mario Zucchelli, showed the presence of Killer Whale *Orcinus orca* (L.), types B and C and Minke Whale (*Balaenoptera bonaerensis* Burmeister). (Lauriano *et al.*, 2007a; 2007b; Lauriano pers.com.). Leopard seals (*Hydrurga leptonyx*) were sighted several times at the end of the slope that penguins climb to reach the colony in the area represented in Map 1.

Environmental characterization

Studies on industrial pollutants in biomarkers allowed to monitor the impact of human activities on the Antarctic biota in Terra Nova Bay area (Focardi *et al.*, 1995; Regoli *et al.*, 1998; Jimenez *et al.*, 1999; Regoli *et al.*, 2005a; Benedetti *et al.*, 2005, 2007; Canapa *et al.*, 2007; Di Bello *et al.*, 2007, Corsolini, 2009).

In Terra Nova Bay, organisms are exposed to a naturally elevated bioavailability of cadmium causing tissue concentrations generally 10–50 folds higher than those typical of temperate species (Mauri *et al.*, 1990; Nigro *et al.*, 1992, 1997; Canapa *et al.*, 2007, Mangano *et al.*, 2014, Caruso *et al.*, 2018). Elevated level of cadmium at Terra Nova Bay modulates bioaccumulation and metabolism of polycyclic aromatic hydrocarbons and of organochlorine xenobiotics in local marine organisms (Regoli *et al.*, 2005a; Benedetti *et al.*, 2007; Canapa *et al.*, 2007). Recent analyses (Signa *et al.* 2019) reported increased concentrations of Pb and Hg (Pb: Grotti *et al.*, 2008; Ianni *et al.*, 2010; Hg: Bargagli *et al.*, 1998; Negri *et al.*, 2006), and phytoplankton reached trace elements levels from 2-fold (Hg) to 4-fold (Cd) and even 10-fold (Pb) higher than those previously recorded (Bargagli *et al.*, 1996, 1998; Dalla Riva *et al.*, 2003). In contrast, Hg concentration measured in feathers of Adélie penguins (*Pygoscelies adeliae*) and Skua (*Catharacta maccormicki*) in 2013 (Signa *et al.* 2019) did not differ from those measured in 1989–1991 (Bargagli *et al.* 1998).

A systematic publication of faunal check-lists for the Terra Nova Bay area has been stated by the Italian National Antarctic Museum (MNA, <https://steu.shinyapps.io/MNA-generale/>) in 2013, with the final target to provide to GBIF distributional information for all taxa occurring in the area. Data are available for: Mollusca (Ghiglione *et al.*, 2013), Tanaidacea (Piazza *et al.*, 2014), Ophiuroidea (Cecchetto *et al.*, 2017), Porifera (Ghiglione *et al.*, 2018), Bryozoa (Cecchetto *et al.*, 2019).

In recent years, remotely operated vehicle surveys and transects were performed. Georeferenced images were taken at specific points identifying the appearing species (Canese *et al.*, 2015). These activities allow to monitor changes in coastal benthic communities (Piazza *et al.*, 2018; Piazza *et al.*, in press). Ongoing studies on food web structure will enable to quantify trophic interactions between species and potential community vulnerability to biodiversity loss and changes in sea-ice dynamics (Calizza *et al.*, 2018, Signa *et al.*, 2019).

Human Activities

The Area is close to the Italian Station Mario Zucchelli (74°41'39"S, 164°06'55"E) that can accommodate approximately 90 people, has facilities for helicopter operations and a jetty for the docking of small boats. Fuel used at the station is Jet-A1. The station is equipped with a waste water treatment plant. Treated water is discharged into the sea adjacent to the station 2.3 km from the northern boundary of the Area. A support ship regularly visits Mario Zucchelli Station during the summer, and there are occasional visits by tourist ships. These usually stop offshore several kilometers to the north of the Area.

Other nearby stations are Gondwana (74°38'0.7"S, 164°13'19" E; Germany), a summer station with capacity for approximately 25 personnel, Jang Bogo station (74°37'15"S, 164°11'57"E; Republic of Korea) year round station with a complement of 60 personnel during summer and 17 during winter. China is currently establishing a new station on nearby Inexpressible Island which will operate year-round with a complement of up to 30 in winter and 80 summer personnel (CAA 2018).

A gravel runway is under construction at Boulder Clay site, Terra Nova Bay (74°44'45"S, 164°01'17"E, 205 m a.s.l.). The end of the runway is about 1.8 km from the penguin colony of Adélie Cove. An Environmental Impact Monitoring Plan has been developed to evaluate changes in the ecosystem during construction and operation of the runway (Draft CEE – MZS gravel runway ATCM39).



6(ii) Access to the Area

Access into the Area is generally by ship. Access into the Area may be made by air or over sea ice when conditions allow. Access routes within the Area have not been defined.

6(iii) Location of structures within and adjacent to the Area

There are no structures within the Area. The nearest structure is the atmospheric monitoring facility (locally referred to as 'Campo Icaro') 650 m north of the northern boundary of the Area, while Mario Zucchelli Station (74°41'42"S, 164°07'23"E) is situated on a small peninsula on the coast adjacent to Tethys Bay, a further 1.65 km to the north. A gravel runway is under construction at Boulder Clay site, Terra Nova Bay (74°44'45"S, 164°01'17"E, 205 m a.s.l.). The end of the runway is about 1.8 km from the penguin colony of Adelie Cove.

6(iv) Location of other protected areas in the vicinity

ASPA No. 175 the high altitude geothermal sites on Mount Melbourne, is a terrestrial site situated 45 km to the NE, which is the only other protected area within close proximity.

6(v) Special zones within the Area

There are no special zones within the Area.

7. Terms and conditions for entry permits

7(i) General permit conditions

Entry into the Area is prohibited except in accordance with a permit issued by the appropriate national authority. Conditions for issuing a permit are that:

- it is issued for scientific purposes, or for educational purposes which cannot be served elsewhere; and/or
- it is issued for essential management purposes consistent with plan objectives such as inspection, maintenance or review;
- the actions permitted will not jeopardise the values of the Area;
- any management activities are in support of the objectives of the Management Plan;
- the actions permitted are in accordance with the Management Plan;
- The permit, or a copy, shall be carried by the holder within the Area;
- permits shall be issued for a stated period

7(ii) Access to and movement within the Area

Access into the Area shall be by sea, land, over sea ice or by air. There are no specific restrictions on routes of access to and movement within the Area, although movements should be kept to the minimum necessary consistent with the objectives of any permitted activities and every reasonable effort should be made to minimise disturbance. Anchoring is prohibited within the Area. There are no overflight restrictions within the Area and aircraft may land by permit when sea ice conditions allow, taking into consideration the Penguin colony situated at Adelie Cove and following the Guidelines for Operations of Aircraft near Concentration of Birds in Antarctica (Resolution 2, 2004), to limit disturbance.

7(iii) Activities that may be conducted within the Area

Activities that may be conducted in the Area should not jeopardise the values of the Area and include:

- Scientific research that cannot be served elsewhere;
- Sampling, which should be the minimum required to reach the scientific goals. Selective and less-invasive sampling methods should always be considered to reduce disturbance of the rich bottom communities;
- Essential management activities, including monitoring and inspection;
- Operational activities in support of scientific research or management of the Area;
- Activities for educational and outreach purposes

7(iv) Installation, modification or removal of structures

Structures or scientific equipment shall not be installed within the Area except as specified in a permit. All markers, structures or scientific equipment installed in the Area shall be clearly identified by country, name of the principal investigator and year of installation. All such items should be made of materials that pose minimal risk of contamination of the Area. Removal of specific equipment for which the permit has expired is mandatory.



7(v) Location of field camps

None within the Area.

7(vi) Restrictions on materials and organisms which can be brought into the Area

- No living animals, plant material, pathogens or microorganisms shall be deliberately introduced into the Area.
- Poultry products, including food products containing uncooked dried eggs, shall not be introduced into the Area.
- No herbicides or pesticides shall be introduced into the Area.
- Chemicals, including radio-nuclides or stable isotopes, which may be introduced for the scientific or management purposes specified in the permit, shall be used in the minimum quantities necessary to achieve the purpose of the activity for which the permit was granted.
- All materials introduced in the Area shall be stored and handled so that risk of their accidental release into the environment is minimized and removed at the end of the period allowed in the permit.
- Visitors shall take special precautions against marine pollution and ensure that sampling equipment or markers brought into the Area are clean. Vessels that are found to show fuel leakage, or a significant risk of such leakage, are prohibited from entering the Area.

7(vii) Taking or harmful interference with native flora or fauna

Taking or harmful interference with native flora or fauna is prohibited, except by permit issued in accordance with Annex II to the Protocol on Environmental Protection to the Antarctic Treaty. Careful environmental evaluation is needed concerning trawling, dragging, grabbing, dredging, or deployment of nets because of the sensitivity of the rich bottom communities to disturbance. More selective and less-invasive sampling methods should always be considered;

Where taking of or harmful interference with animals is involved, the *SCAR Code of Conduct for the Use of Animals for Scientific Purposes in Antarctica* (ATCM XXXIV-CEP XIV, 2011) should be used as a minimum standard.

7(viii) Collection or removal of materials not brought into the Area by the permit holder

Any anthropogenic material found should be notified to the appropriate national authority.

Material may be collected or removed from the Area only in accordance with a permit. In this case removal of material should not create an impact greater than leaving the material *in situ*.

7(ix) Disposal of waste

All wastes, including all human wastes, shall be removed from the Area.

7(x) Measures that may be necessary to continue to meet the aims of the Management Plan

Permits may be granted to enter the Area to

- carry out monitoring and site inspection activities, which may involve the collection of limited samples for analysis or review, or for protective measures;
- Install markers on specific sites of long-term monitoring.

7(xi) Requirements for reports

The holder of each permit issued should report to the appropriate national competent authority about the activity undertaken in the Area.

Such reports should include, as appropriate, the information identified in Appendix 2- ASPA visit report form of the Guide to the Preparation of Management plans for ASPAs (Resolution 2 , 2011). Parties should, wherever possible, exchange with the Party that proposed the Management Plan, information on reports received to assist managing the Area.



8. References

- Accornero A., Manno C., Arrigo K.R., Martini Atucci S., 2003. "The vertical flux of particulate matter in the polynya of Terra Nova Bay. Part I. Chemical constituents" *Antarctic Science* 15 (1), 119-132.
- Albertelli G., Cattaneo-Vietti R., Chiantore M., Pusceddu A., Fabiano M., 1998. Food availability to an Adamussium bed during the austral Summer 1993/94 (Terra Nova Bay, Ross Sea). *Journal of Marine Systems* 17: 425-34.
- Alvaro M.C, Blazewicz-Paszkowycz M., Davey N., Schiaparelli S., 2011. Skin-digging tanaidaceans: the unusual parasitic behaviour of *Exspina typica* (Lang, 1968) in Antarctic waters and worldwide deep basins. *Antarct Sci*, vol. 23 (4); p. 343-348, ISSN: 0954-1020, doi: 10.1017/S0954102011000186
- Ansell A.D., Cattaneo-Vietti R., Chiantore M., 1998. Swimming in the Antarctic scallop *Adamussium colbecki*: analysis of in situ video recordings. *Antarctic Science* 10 (4): 369-75.
- Ballerini T., Tavecchia G., Olmastroni S., Pezzo F., Focardi S., 2009. Nonlinear effects of winter sea ice on the survival probabilities of Adélie penguins. *Oecologia* 161:253-265.
- Bargagli R., Nelli L., Ancora S., Focardi S., 1996. Elevated cadmium accumulation in marine organisms from Terra Nova bay (Antarctica). *Polar Biology* 16: 513-520.
- Bargagli R., Monaci F., Sanchez-Hernandez J.C., Cateni D., 1998. Biomagnification of mercury in an Antarctic marine coastal food web. *Marine Ecology Progress Series* 169: 65-76.
- Bargagli R., 2005. *Antarctic Ecosystems. Environmental Contamination, Climate Change, and Human Impact. Ecological Studies*, vol. 175; Springer-Verlag, Heidelberg, 395 pp.
- Bargagli R., 2008. Environmental contamination in Antarctic ecosystems. *Sci. Total Environ.* 400: 212-226.
- Bavestrello G., Arillo A., Calcinaï B., Cattaneo-Vietti R., Cerrano C., Gaino E., Penna A., Sara' M., 2000. Parasitic diatoms inside Antarctic sponges. *Biol. Bull.* 198: 29-33.
- Benedetti M., Gorbi S., Bocchetti R., Fattorini D., Notti A., Martuccio G., Nigro M., Regoli F. (2005). Characterization of cytochrome P450 in the Antarctic key sentinel species *Trematomus bernacchii*. *Pharmacologyonline* 3: 1-8 ISSN-1827-8620.
- Benedetti M., Martuccio G., Fattorini D., Canapa A., Barucca M., Nigro M., Regoli F. (2007). Oxidative and modulatory effects of trace metals on metabolism of polycyclic aromatic hydrocarbons in the Antarctic fish *Trematomus bernacchii*. *Aquat. Toxicol.* 85: 167-175
- Berkman P.A., Nigro M., 1992. Trace metal concentrations in scallops around Antarctica: Extending the Mussel Watch Programme to the Southern Ocean. *Marine Pollution Bulletin* 24 (124): 322-23.
- Borghesi N., Corsolini S., Focardi S., 2008. Levels of polybrominated diphenyl ethers (PBDEs) and organochlorine pollutants in two species of Antarctic fish (*Chionodraco hamatus* and *Trematomus bernacchii*). *Chemosphere*, 73, 155-160.
- Bruni V., Maugeri M.L., Monticelli L.S., 1997. Faecal pollution indicators in the Terra Nova Bay (Ross Sea, Antarctica). *Marine Pollution Bulletin* 34 (11): 908-12.
- Budillon G., Spezie G., 2000. "Thermoaline structure and variability in the Terra Nova Bay polynya (Ross Sea) between 1995-98". *Antarctic science* 12, 243-254.
- Calizza E., Careddu G., Sporta Caputi S., Rossi L., Costantini M.L., 2018. Time- and depth-wise trophic niche shifts in Antarctic benthos. *PLoS one* 13: e0194796.
- Canapa A., Barucca M., Gorbi S., Benedetti M., Zucchi S., Biscotti M.A., Olmo E., Nigro M., Regoli F., 2007. Vitellogenin gene expression in males of the Antarctic fish *Trematomus bernacchii* from Terra Nova Bay (Ross Sea): A role for environmental cadmium? *Chemosphere*, 66:1270-1277.
- Canese S., Mazzoli C., Montagna P., Schiaparelli S., Taviani M., 2015. The Terra Nova Bay 'Canyon': ROV survey of nearshore shallow to deep carbonate factories. XII International Symposium on Antarctic Earth Sciences ISAES, 13-17 July 2015, Goa, India.
- Cantone G., Castelli A., Gambi M.C., 2000. The Polychaete fauna off Terra Nova Bay and Ross Sea: biogeography, structural aspects and ecological role. In: *Ross Sea Ecology*, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 551-61.
- Caruso C., Rizzo C., Mangano S., Poli A., Di Donato P., Nicolaus B., Di Marco G., Michaud L., Lo Giudice A. (2018). Extracellular polymeric substances with metal adsorption capacity produced by *Pseudoalteromonas* sp. MER144 from Antarctic seawater. *Environmental Science and Pollution Research*, 25: 4667-4677.
- Castellano M. 2006. "Aspetti trofo-funzionali dell'ecosistema marino costiero antartico: sostanza organica particellata e disciolta", Univeristà degli Studi di Genova, PhD Thesys.
- Cattaneo-Vietti R., 1991. Nudibranch Molluscs from the Ross Sea, Antarctica. *J. Moll. Stud.* 57: 223-28.
- Cattaneo-Vietti R., Bavestrello G., Cerrano C., Gaino E., Mazzella L., Pansini M., Sarà M., 2000c. The role of sponges of Terra Nova Bay ecosystem. In: *Ross Sea Ecology*, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 539-49.



- Cattaneo-Vietti R., Bavestrello G., Cerrano C., Sara' M., Benatti U., Giovine M., Gaino E., 1996. Optical fibres in an Antarctic sponge. *Nature* 383: 397-98.
- Cattaneo-Vietti R., Chiantore M., Albertelli G., 1997. The population structure and ecology of the Antarctic Scallop, *Adamussium colbecki* in Terra Nova Bay (Ross Sea, Antarctica). *Scientia Marina* 61 (Suppl. 2): 15-24.
- Cattaneo-Vietti R., Chiantore M., Gambi M.C., Albertelli G., Cormaci M., Di Geronimo I., 2000a. Spatial and vertical distribution of benthic littoral communities in Terra Nova Bay. In: Ross Sea Ecology, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 503-14.
- Cattaneo-Vietti R., Chiantore M., Mistic C., Povero P., Fabiano M., 1999. The role of pelagic-benthic coupling in structuring littoral benthic communities at Terra Nova Bay (Ross Sea) and inside the Strait of Magellan. *Scientia Marina* 63 (Supl. 1): 113-21.
- Cattaneo-Vietti R., Chiantore M., Schiaparelli S., Albertelli G., 2000b. Shallow and deep-water mollusc distribution at Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology* 23: 173-82.
- Cecchetto M., Alvaro M.C., Ghiglione C., Guzzi A., Mazzoli C., Piazza P., Schiaparelli S., 2017. Distributional records of Antarctic and sub-Antarctic Ophiuroidea from samples curated at the Italian National Antarctic Museum (MNA): check-list update of the group in the Terra Nova Bay area (Ross Sea) and launch of the MNA 3D model 'virtual gallery'. *ZooKeys*, 705: 61-79. <https://doi.org/10.3897/zookeys.705.13712>.
- Cecchetto M., Lombardi C., Canese S., Cocito S., Kuklinski P., Mazzoli C., Schiaparelli S., 2019. Bryozoa collection of the Italian National Antarctic Museum (MNA), with an updated checklist from Terra Nova Bay (Ross Sea). *Zookeys* 812: 1-22.
- Cerrano C., Calcinai B., Cucchiari E., Di Camillo C., Nigro M., Regoli F., Sarà A., Schiaparelli S., Totti C., Bavestrello G., 2004. Are diatoms a food source for Antarctic sponges?. *Chemistry and Ecology*, vol. 20: 57-64.
- Cerrano C., Arillo A., Bavestrello G., Calcinai B., Cattaneo-Vietti R., Penna A., Sarà M., Totti C., 2000a. Diatom invasion in the Antarctic hexactinellid sponge *Scolymastra joubini*. *Polar Biology* 23: 441-44.
- Cerrano C., Bavestrello G., Calcinai B., Cattaneo-Vietti R., Sarà A., 2000b. Asteroids eating sponges from Tethys Bay, East Antarctica. *Antarctic Science* 12(4): 431-32.
- Cerrano C., G. Bavestrello, B. Calcinai, R. Cattaneo-Vietti, M. Chiantore, M. Guidetti, A. Sarà, 2001a. Bioerosive processes in Antarctic seas. *Polar Biology* 24: 790-92.
- Cerrano C., Puce S., Chiantore M., Bavestrello G., 2000c. Unusual trophic strategies of *Hydractinia angusta* (Cnidaria, Hydrozoa) from Terra Nova Bay, Antarctica. *Polar Biology* 23(7): 488-94.
- Cerrano C., S. Puce, M. Chiantore, G. Bavestrello, R. Cattaneo-Vietti, 2001b. The influence of the epizooic hydroid *Hydractinia angusta* on the recruitment of the Antarctic scallop *Adamussium colbecki*. *Polar Biology* 24: 577-81.
- Chiantore M., Cattaneo-Vietti R., Albertelli G., Mistic M., Fabiano M., 1998. Role of filtering and biodeposition by *Adamussium colbecki* in circulation of organic matter in Terra Nova Bay (Ross Sea, Antarctica). *Journal of Marine Systems* 17: 411-24.
- Chiantore M., Cattaneo-Vietti R., Berkman P.A., Nigro M., Vacchi M., Schiaparelli S., Albertelli G., 2001. Antarctic scallop (*Adamussium colbecki*) spatial population variability along the Victoria Land Coast, Antarctica. *Polar Biology* 24: 139-43.
- Chiantore M., Cattaneo-Vietti R., Povero P., Albertelli G., 2000. The population structure and ecology of the antarctic scallop *Adamussium colbecki* in Terra Nova Bay. In: Ross Sea Ecology, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 563-73.
- Chiantore M., R. Cattaneo-Vietti, L. Elia, M. Guidetti, M. Antonini, 2002. Reproduction and condition of the scallop *Adamussium colbecki* (Smith 1902), the sea-urchin *Sterechinus neumayeri* (Meissner, 1900) and the sea-star *Odontaster validus* Koehler, 1911 at Terra Nova Bay (Ross Sea): different strategies related to inter-annual variations in food availability. *Polar Biology* 22: 251-55.
- Chiantore M., Vacchi M., 2012. Dense populations of the Antarctic scallop *Adamussium colbecki* in Terra Nova Bay (Subarea 88.1J): potential VMEs adjacent to the Terra Nova Bay ASPA (No. 161). CCAMLR WG-MME-12/23, 12 pp.
- Cormaci M., Furnari G., Scammacca B., 1992b. The benthic algal flora of Terra Nova Bay (Ross Sea, Antarctica). *Botanica Marina* 35(6): 541-52
- Cormaci M., Furnari G., Scammacca B., 1992c. Carta della vegetazione marina di Baia Terra Nova (Mare di Ross, Antartide). *Biologia Marina* 1: 313-14.
- Cormaci M., Furnari G., Scammacca B., Alongi G., 1996. Summer biomass of a population of *Iridaea cordata* (Gigartinaceae, Rhodophyta) from Antarctica. In: Lindstrom SC, Chapman DJ (Eds) Proceedings of the XV Seaweeds Symposium. *Hydrobiologia* 326/327: 267-72.
- Cormaci M., Furnari G., Scammacca B., Casazza G., 1992a. Il fitobenthos di Baia Terra Nova (Mare di Ross, Antartide): osservazioni sulla flora e sulla zonazione dei popolamenti. In: Gallardo VA, Ferretti O, Moyano HI (eds) Actas del Semin. Int. Oceanografía in Antartide. Centro EULA, Universidad de Concepción, Chile. ENEA: 395-408.
- Corsolini S, Nigro M, Olmastroni S, Focardi S, Regoli F 2001 Susceptibility to oxidative stress in Adelie and Emperor penguin, *Polar Biology*, vol. 24: 365-368.
- Corsolini S, Borghesi N., Ademolo N., Focardi S., 2011. Chlorinated biphenyls and pesticides in migrating and resident seabirds from East and West Antarctica. *Environment International* 37(8): 1329-1335.



- Corsolini S., 2009. Industrial contaminants in Antarctic biota. *Journal of Chromatography A*, 1216, 598–612.
- Corsolini S., 2011. Antarctic: Persistent Organic Pollutants and Environmental Health in the Region. In: Nriagu JO (ed.) *Encyclopedia of Environmental Health*, volume 1, pp. 83–96 Burlington: Elsevier, NVRN/978-0-444-52273-3.
- Corsolini S., Kannan K., Imagawa T., Focardi S., Giesy J.P., 2002. Polychloronaphthalenes and other dioxin-like compounds in Arctic and Antarctic marine food webs. *Environmental Science and Technology*, 36: 3490–3496.
- Dalla Riva S., Abelloschi M.L., Magi E., Soggia F., 2004. The utilization of the antarctic environmental specimen bank (BCAA) in monitoring Cd and Hg in an antarctic coastal area in Terra Nova bay (Ross Sea – Northern Victoria land). *Chemosphere* 56: 59–69.
- Di Bello D., Vaccaio E., Longo V., Regoli F., Nigro M., Benedetti M., Gervasi PG, Pretti C. (2007). Presence and inducibility by β -Naphthoflavone of CYP 1A1, CYP 1B1, UDP-GT, GST and DT-Diaphorase enzymes in *Trematomus bernacchii*, an Antarctic fish. *Aquatic Toxicol.* 84: 19–26
- Fabiano M., Chiantore M., Povero P., Cattaneo-Vietti R., Pusceddu A., Misisic C., Albertelli G., 1997. Short-term variations in particulate matter flux in Terra Nova Bay, Ross Sea. *Antarctic Science* 9(2): 143–149.
- Fabiano M., Danovaro R., Crisafi E., La Ferla R., Povero P., Acosta Pomar L., 1995. Particulate matter composition and bacterial distribution in Terra Nova Bay (Antarctica) during summer 1989–90. *Polar Biology* 15: 393–400.
- Fabiano M., Povero P., Danovaro R., 1996. Particulate organic matter composition in Terra Nova Bay (Ross Sea, Antarctica) during summer 1990. *Antarctic Science* 8(1): 7–13.
- Focardi S., Fossi M.C., Lari L., Casini S., Leonzio C., Meidel S.K., Nigro M., 1995. Induction of MFO Activity in the Antarctic fish *Pagothenia bernacchii*: Preliminary results. *Marine Environmental Research.*, 39: 97–100.
- Focardi S., Bargagli R., Corsolini S., 1993. Organochlorines in marine Antarctic food chain at Terra Nova Bay (Ross Sea). *Korean Journal of Polar Research* 4: 73–77.
- Frezzotti, M., Salvatore, M.C., Vittuari, L., Grigioni, P., De Silvestri L., 2001. Satellite Image Map: Northern Foothills and Inexpressible Island Area (Victoria Land, Antarctica). *Terra Antarctica Reports* n° 6, 8 p. + map – ISBN 88-900221-9-1.
- Gaino E., Bavestrello G., Cattaneo-Vietti R., Sara' M., 1994. Scanning electron microscope evidence for diatom uptake by two Antarctic sponges. *Polar Biology* 14: 55–58.
- Gambi M.C., Buia M.C., Mazzella L., Lorenti M., Scipione M.B., 2000a. Spatio-temporal variability in the structure of benthic populations in a physically controlled system off Terra Nova Bay: the shallow hard bottoms. In: *Ross Sea Ecology*, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 527–538.
- Gambi M.C., Castelli A., Guizzardi M., 1997. Polychaete populations of the shallow soft bottoms off Terra Nova Bay (Ross Sea, Antarctica): distribution, diversity and biomass. *Polar Biology* 17: 199–210.
- Gambi M.C., Giangrande A., Patti F.P., 2000b. Comparative observations on reproductive biology of four species of *Perkinsiana* (Polychaeta, Sabellidae). *Bulletin of Marine Science* 67(1): 299–309.
- Gambi M.C., Lorenti M., Russo G.F., Scipione M.B., 1994. Benthic associations of the shallow hard bottoms off Terra Nova Bay (Ross Sea, Antarctica): zonation, biomass and population structure. *Antarctic Science* 6(4): 449–62.
- Gavagnin M., Trivellone E., Castelluccio F., Cimino G., Cattaneo-Vietti R., 1995. Glyceryl ester of a new halimane diterpenoic acid from the skin of the antarctic nudibranch *Austrodoris kerguelensis*. *Tetrahedron Letters* 36: 7319–22.
- Ghiglione C, Alvaro M.C., Griffiths H.J., Linse K., Schiaparelli S., 2013. Ross Sea Mollusca from the Latitudinal Gradient Program: R/V *Italica* 2004 Rauschert dredge samples. *ZooKeys*, 341: 37–48.
- Grotti M., Soggia F., Lagomarsino C., Dalla Riva S., Goessler W., Francesconi, K.A., 2008. Natural variability and distribution of trace elements in marine organisms from Antarctic coastal environments. *Antarctic Science* 20: 39–51.
- Guglielmo G., Zagami G., Saggiorno V., Catalano G., Granata A., 2007. "Copepods in spring annual sea ice at Terra Nova Bay (Ross Sea, Antarctica)" *Polar Biology* 30, 747–758.
- Guglielmo L., Carrada G.C., Catalano G., Dell'Anno A., Fabiano M., Lazzara L., Mangoni O., Pusceddu A., Saggiorno V., 2000. Structural and functional properties of sympagic communities in the annual sea ice at Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology* 23(2): 137–46.
- Guglielmo L., Granata A., Greco S., 1998. Distribution and abundance of postlarval and juvenile *Pleuragramma antarcticum* (Pisces, Nototheniidae) of Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology* 19: 37–51.
- Humphries G.R.W., Che-Castaldo C., Naveen R., Schwaller M., McDowall P., Schrimpf M., and Lynch H.J. 2017. Mapping Application for Penguin Populations and Projected Dynamics (MAPPPD): Data and tools for dynamic management and decision support. *Polar Records* .
- Ianni C., Magi E., Soggia F., Rivaro P., Frache R., 2010. Trace metal speciation in coastal and off-shore sediments from Ross Sea (Antarctica). *Microchemical Journal* 96: 203–212
- Jimenez B., Fossi M.C., Nigro M., Focardi S., 1999. Biomarker approach to evaluating the impact of scientific stations on the Antarctic environment using *trematomus bernacchii* as a bioindicator organism. *Chemosphere*, 39: 2073–2078.



- La Mesa M., Arneri E., Giannetti G., Greco S., Vacchi M., 1996. Age and growth of the nototheniid fish *Trematomus bernacchii* Boulenger from Terra Nova Bay, Antarctica. *Polar Biology* 16: 139-45.
- La Mesa M., J.T. Eastman, M. Vacchi, 2004. The role of nototheniid fish in the food web of the Ross Sea shelf waters: a review. *Polar Biol.*, 27: 321-338.
- La Mesa M., Vacchi M., Castelli A., Diviacco G., 1997. Feeding ecology of two nototheniid fishes *Trematomus hansonii* and *Trematomus loennbergi* from Terra Nova Bay, Ross Sea. *Polar Biology* 17: 62-68.
- La Mesa M., Vacchi M., T. Zunini Sertorio, 2000. Feeding plasticity of *Trematomus newnesi* (Pisces, Nototheniidae) in Terra Nova Bay, Ross Sea, in relation to environmental conditions. *Polar Biology* 23(1): 38-45.
- Lauriano G., Fortuna C.M., Vacchi M., 2007a. Observation of killer whale (*Orcinus orca*)
- Lauriano G., Vacchi M., Ainley D., Ballard G., 2007b. Observations of top predators foraging on fish in the pack ice of the southern Ross Sea. *Antarctic Science*, 19(4): 439-440.
- Lo Giudice A., Casella P., Bruni V., Michaud L. (2013). Response of bacterial isolates from Antarctic shallow sediments towards heavy metals, antibiotics and polychlorinated biphenyls. *Ecotoxicology*, 22: 240-250
- Lo Giudice A., Azzaro M., Schiaparelli S. (2019). Microbial Symbionts of Antarctic Marine Benthic Invertebrates. In *The Ecological Role of Micro-organisms in the Antarctic Environment*. Castro-Sowinski S. (Ed.), Chapter 13, Springer Polar Sciences. Pp. 277-296. https://doi.org/10.1007/978-3-030-02786-5_13.
- Mangoni O., Modigh M., Conversano F., Carrada G.C., Saggiorno V., 2004. "Effects of summer ice coverage on phytoplankton assemblages in the Ross Sea, Antarctica" *Deep-Sea Research I*, 51, 1601-1617.
- Massolo S., Messa R., Rivaro P., Leardi R., 2009. "Annual and spatial variations of chemical and physical properties in the Ross Sea surface waters (Antarctica)" *Continental Shel Research* 29, 2333-2344.
- Mauri M., Orlando E., Nigro M., Regoli F., 1990. Heavy metals in the Antarctic scallop *Adamussium colbecki* (Smith). *Mar. Ecol. Progr. Ser.* 67: 27-33.
- Minganti V., Capelli R., Fiorentino F., De Pellegrini R., Vacchi M., 1995. Variations of mercury and selenium concentrations in *Adamussium colbecki* and *Pagothenia bernacchii* from Terra Nova Bay (Antarctica) during a five year period. *Int. J. Environ. Anal. Chem.* 61: 239-48.
- Negri A., Burns K., Boyle S., Brinkman D., Webster N., 2006. Contamination in sediments, bivalves and sponges of McMurdo Sound, Antarctica. *Environmental Pollution* 143: 456-467.
- Nigro M., Orlando E., Regoli F., 1992. Ultrastructural localisation of metal binding sites in the kidney of the Antarctic scallop *Adamussium colbecki*. *Marine Biology*, 113: 637-643.
- Nigro M., Regoli F., Rocchi R., Orlando E., 1997. Heavy metals in Antarctic Molluscs. In "Antarctic Communities" (B. Battaglia, J. Valencia and D.W.H Walton Eds.), Cambridge University Press, 409-412
- Nonnis Marzano F., Fiori F., Jia G., Chiantore M., 2000. Anthropogenic radionuclides bioaccumulation in Antarctic marine fauna and its ecological relevance. *Polar Biology* 23: 753-58.
- Pane L., Feletti M., Francomacaro B., Mariottini G.L., 2004. "Summer coastal zooplankton biomass and copepod community structure near the Italian Terra Nova Base (Terra Nova Bay, Ross Sea, Antarctica)" *Journal of Plankton Research*, vol 26, issue 12, 1479-1488.
- Piazza et al (IN PRESS) Antarctic benthos monitoring using underwater photogrammetry: lessons learned, legacy data rescue and future developments for long-term monitoring programmes. *Polar Biology* (POBI-D-18-00100).
- Piazza P., Cummings V., Lohrer D., Marini S., Marriott P., Menna F., Nocerino E., Peirano A., Schiaparelli S., 2018. Divers-operated underwater photogrammetry: applications in the study of Antarctic benthos. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, Volume XLII-2, 885-892. <https://doi.org/10.5194/isprs-archives-XLII-2-885-2018>.
- Piazza P., Błazewicz-Paszkowycz M., Ghiglione C., Alvaro M.C., Schnabel K., Schiaparelli S., 2014. Distributional records of Ross Sea (Antarctica) Tanaidacea from museum samples stored in the collections of the Italian National Antarctic Museum (MNA) and the New Zealand National Institute of Water and Atmospheric Research (NIWA). *ZooKeys*, 451: 49-60.
- Povero P., Castellano M., Ruggieri N., Monticelli L.S., Saggiorno V., Chiantore M.C., Guidetti M., Cattaneo-Vietti R., 2006. "Water column features and their relationship with sediments and benthic communities along the Victoria Land coast, Ross Sea, Antarctica, summer 2004" *Antarctic Science* 18 (4), 603-613.
- Povero P., Chiantore M., Mistic C., Budillon G., Cattaneo-Vietti R., 2001. Pelagic-benthic coupling in Adélie Cove (Terra Nova Bay, Antarctica): a strongly land forcing controlled system? *Polar Biology* 24: 875-882.
- Puce S., Cerrano C., Bavestrello G., 2002. *Eudendrium* (Cnidaria, Anthomedusae) from the Antarctic Ocean with a description of new species. *Polar Biology* 25: 366-73.
- Pusceddu A., Cattaneo-Vietti R., Albertelli G., Fabiano M., 1999. Origin, biochemical composition and vertical flux of particulate organic matter under the pack ice in Terra Nova Bay (Ross Sea, Antarctica) during late summer 1995. *Polar Biology* 22: 124-32.



- Regoli F., Nigro M., Benedetti M., Fattorini D., Gorbi S., 2005b. Antioxidant efficiency in early life stages of the Antarctic silverfish *Pleuragramma antarcticum*: Responsiveness to pro-oxidant conditions of platelet ice and chemical exposure. *Aquatic Toxicology*, vol. 75: 43-52.
- Regoli F., Nigro M., Benedetti M., Gorbi S., Pretti C., Gervasi P.G., Fattorini D., 2005a. Interactions between metabolism of trace metals and xenobiotics agonist of the aryl hydrocarbon receptor in the Antarctic fish *Trematomus bernacchii*: environmental perspectives. *Environmental Toxicology and Chemistry*, vol. 24(6): 201-208
- Regoli F., Nigro M., Bertoli E., Principato G.B., Orlando E., 1997b. Defences against oxidative stress in the Antarctic scallop *Adamussium colbecki* and effects of acute exposure to metals. *Hydrobiologia*, 355: 139-144.
- Regoli F., Nigro M., Bompadre S., Wiston G., 2000a. Total oxidant scavenging capacity (TOSC) of microsomal and cytosolic fractions from Antarctic Arctic and Mediterranean Scallops: differentiation between three different potent oxidants. *Aquatic Toxicology*, 49: 13-25.
- Regoli F., Nigro M., Chiantore M.C., Gorbi S., Wiston G., 2000b. Total oxidant scavenging capacity of Antarctic, Arctic and Mediterranean scallops. *Italian Journal of Zoology*, vol. 67: 5-94.
- Regoli F., Nigro M., Chierici E., Cerrano C., Schiaparelli S., Totti C., Bavestrello G., 2004. Variations of antioxidant efficiency and presence of endosymbiotic diatoms in the Antarctic porifera *Haliclona dancoi*. *Marine Environmental Research*, vol. 58: 637-640.
- Regoli F., Nigro M., Orlando E., 1998. Lysosomal and antioxidant defences to metals in the Antarctic scallop *Adamussium colbecki*. *Aquatic Toxicology*, 40: 375-392.
- Regoli F., Principato G.B., Bertoli E., Nigro M., Orlando E., 1997a. Biochemical characterisation of the antioxidant system in the scallop *Adamussium colbecki*, a sentinel organism for monitoring the Antarctic environment. *Polar Biology*, 17: 251-25.
- Regoli F., M. Nigro, M. Chiantore, G.W. Winston, 2002. Seasonal variations of susceptibility to oxidative stress in *Adamussium colbecki*, a key bioindicator species for the Antarctic marine environment. *The Science of the Total Environment*, 289: 205-211.
- Regoli F., Nigro M., Chierici E., Cerrano C., Schiaparelli S., Totti C., Bavestrello G., 2004. Variations of antioxidant efficiency and presence of endosymbiotic diatoms in the Antarctic porifera *Haliclona dancoi*. *Marine Environmental Research*, 58: 637-640.
- Sarà A., Cerrano C., Sarà M., 2002. Viviparous development in the Antarctic sponge *Stylocordyla borealis* Loven, 1868. *Polar Biology* 25: 425-31.
- Sarà M., Balduzzi A., Barbieri M., Bavestrello G., Burlando B., 1992. Biogeographic traits and checklist of Antarctic demosponges. *Polar Biology* 12: 559-85.
- Schiaparelli S., Aliani, 2019. Oceanographic moorings as year-round laboratories for investigating growth performance and settlement dynamics in the Antarctic scallop *Adamussium colbecki* (E.A. Smith, 1902). *PeerJ* 7:e6373, DOI 10.7717/peerj.6373
- Schiaparelli S., Linse K., 2006. A reassessment of the distribution of the common Antarctic scallop *Adamussium colbecki* (Smith, 1902). *Deep-Sea Research II*, 53: 912-920.
- Schiaparelli S., Albertelli G., Cattaneo-Vietti R., 2003. The epibiotic assembly on the sponge *Haliclona dancoi* (Topsent, 1901) at Terra Nova Bay (Antarctica, Ross Sea). *Polar Biology*, 26: 342-347.
- Schiaparelli S., Alvaro M.C., Kilgallen N., Scinto A., Lorz A.N., 2015. Host-shift speciation in Antarctic symbiotic invertebrates: further evidence from the new amphipod species *Lepidepecrella debroyeri* from the Ross Sea? *Hydrobiologia* 761: 143-159.
- Schiaparelli S., Alvaro M.C.; Barnich R., 2011. Polynoid polychaetes living in the gut of irregular sea urchins: a first case of inquilinism in the Southern Ocean and an overview of polychaete-echinoderm associations. *Antarctic Science*, 144- 151 23 (2).
- Schiaparelli S., Cattaneo-Vietti R., Chiantore M., 2000. Adaptive morphology of *Capulus subcompressus* Pelseneer, 1903 (Gastropoda: Capulidae) from Terra Nova Bay, Ross Sea (Antarctica). *Polar Biology* 23: 11-16.
- Schiaparelli S., Ghirardo C., Bohn J., Chiantore M., Albertelli G., Cattaneo-Vietti R. 2007. Antarctic associations: the parasitic relationship between the gastropod *Bathycrinicola tumidula* (Thiele, 1912) (Ptenoglossa: Eulimidae) and the comatulid *Notocrinus virilis* Mortensen, 1917 (Crinoidea: Notocrinidae) in the Ross Sea. *Polar Biology*, 30: 1545-1555.
- Schiaparelli S., Lörz A.N., Cattaneo-Vietti R., 2006. Diversity and distribution of mollusc assemblages on the Victoria Land coast and the Balleny Islands, Ross Sea, Antarctica. *Antarctic Science*, 18 (4): 615-631.
- Schiaparelli S., Cattaneo-Vietti R., Mierzejewski P., 2004. A "protective shell" around the larval cocoon of *Cephalodiscus densus* Andersson, 1907 Graptolithoidea (Hemichordata). *Polar Biology*, 27: 813-817.
- Schiaparelli S., Alvaro M.C., Bohn J., Albertelli G., 2010. "Hitchhiker" polynoid polychaetes in cold deep waters and their potential influence on benthic soft bottom food webs. *Antarctic Science*, 399- 407 22 (4).
- Schiaparelli S., Jirkov I.A., 2016A reassessment of the genus *Amphicteis* Grube, 1850 (Polychaeta: Amphaeidae) with the description of *Amphicteis teresae* sp nov from Terra Nova Bay (Ross Sea, Antarctica). *Italian Journal of Zoology* 83: 531-542.
- Signa G., Calizza E., Costantini, M.L., Tramati C., Sporta Caputi S., Mazzola A., Rossi L., Vizzini, S., 2019. Horizontal and vertical food web structure drives trace element trophic transfer in Terra Nova Bay, Antarctica. *Environmental Pollution* 246: 772-781.
- Simeoni U., Baroni C., Meccheri M., Taviani M., Zanon G., 1989. Coastal studies in Northern Victoria Land (Antarctica): Holocene beaches of Inexpressible island, Tethys Bay and Edmonson Point. *Boll. Ocean. Teor. Appl.* 7(1-2): 5-16.



- Stocchino C., Lusetti C., 1988. Le costanti armoniche di marea di Baia Terra Nova (Mare di Ross, Antartide). F.C. 1128 Istituto Idrografico della Marina, Genova.
- Stocchino C., Lusetti C., 1990. Prime osservazioni sulle caratteristiche idrologiche e dinamiche di Baia Terra Nova (Mare di Ross, Antartide). F.C. 1132 Istituto Idrografico della Marina, Genova.
- Swadling K.M., Penot F., Vallet C., Rouyer A., Gasparini S., Mousseau L., Smith M., Goffart A., Koubbi P., 2003. "Interannual variability of zooplankton in the Dumont d'Urville sea (39°E-146°E), east Antarctica, 2004-2008" *Polar Science* 5, 118-133, (2011)
- Tagliabue A. & Arrigo K.R., "Anomalously low zooplankton abundance in the Ross Sea: An alternative explanation" *Limnol. Oceanogr.* 48, 686-699.
- Vacchi M., Cattaneo-Vietti R., Chiantore M., Dalù M., 2000a. Predator-prey relationship between nototheniid fish *Trematomus bernacchii* and Antarctic scallop *Adamussium colbecki* at Terra Nova Bay (Ross Sea). *Antarctic Science* 12(1): 64-68.
- Vacchi M., Greco S., 1994a. Capture of the giant Nototheniid fish *Dissostichus mawsoni* in Terra Nova Bay (Antarctica): Notes on the fishing equipment and the specimens caught. *Cybium* 18(2): 199-203.
- Vacchi M., Greco S., La Mesa M., 1991. Ichthyological survey by fixed gears in Terra Nova Bay (Antarctica). Fish list and first results. *Memorie di Biologia Marina e di Oceanografia* 19: 197-202.
- Vacchi M., La Mesa M., 1995. The diet of Antarctic fish *Trematomus newnesi* Boulenger, 1902 (Notothenidae) from Terra Nova Bay, Ross Sea. *Antarctic Science* 7(1): 37-38.
- Vacchi M., La Mesa M., 1997. Morphometry of *Cryodraco* specimens of Terra Nova Bay. *Cybium* 21(4): 363-68.
- Vacchi M., La Mesa M., Castelli A., 1994b. Diet of two coastal nototheniid fish from Terra Nova Bay, Ross Sea. *Antarctic Science* 6(1): 61-65.
- Vacchi M., La Mesa M., Greco S., 2000b. The coastal fish fauna of Terra Nova Bay, Ross Sea (Antarctica). In: *Ross Sea Ecology*, F. Faranda, L. Guglielmo and A. Ianora Eds., Springer Verlag, Berlin Heidelberg: 457-68.
- Vacchi M., La Mesa M., Eastman J.T., 2004a. "The role of nototheniid fish in the food web of the Ross Sea shelf waters: a review" *Polar Biology* 27(6), 321-338, (2004)
- Vacchi M., La Mesa M., Dalù M., MacDonald J., 2004b. Early life stages in the life cycle of Antarctic silverfish, *Pleuragramma antarcticum* in Terra Nova Bay, Ross Sea. *Antarctic Science*
- Vacchi M., Romanelli M., La Mesa M., 1992. Age structure of *Chionodraco hamatus* (Teleostei, Channichthyidae) samples caught in Terra Nova Bay, East Antarctica. *Polar Biology* 12: 735-38.
- Van dijk G.L., Arrigo K.R., 2005. "Annual cycles of sea ice and phytoplankton in three Ross Sea polynyas" Poster at 3rd International Conference on the Oceanography of the Ross Sea Antarctica. Venezia, Italy, 10-14 Oct.



Map 1 Terra Nova Bay ASPA N° 161, Victoria Land, Ross Sea.

