



Management Plan

for Antarctic Specially Protected Area No. 110 LYNCH ISLAND, SOUTH ORKNEY ISLANDS

Introduction

The primary reason for the designation of Lynch Island, South Orkney Islands (Latitude 60°39'10" S, Longitude 045°36'25" W; 0.14 km²), as Antarctic Specially Protected Area (ASPA) 110 is to protect environmental values, and primarily the terrestrial flora within the Area.

Lynch Island, Marshal Bay, South Orkney Islands, was originally designated as a Specially Protected Area through Recommendation IV-14 (1966, SPA No. 14) after a proposal by the United Kingdom. It was designated on the grounds that the island "supports one of the most extensive and dense areas of grass (*Deschampsia antarctica*) known in the Treaty area and that it provides an outstanding example of a rare natural ecological system". These values were amplified and extended by Recommendation XVI-6 (1991) when a management plan for the site was adopted.

Lynch Island is 2.4 km from Signy Island, the location of Signy Research Station (UK), and about 200 m from Coronation Island, the largest of the South Orkney Islands. The Area has been afforded special protection for most of the modern era of scientific activity in the region, with entry permits having been issued only for compelling scientific reasons. Thus, the island has not been subjected to frequent visits, scientific research or sampling. Since 1983, the numbers of Antarctic fur seals in the South Orkney Islands has increased significantly, with consequent destruction of accessible areas of vegetation where the seals come ashore. Some vegetated areas on Lynch Island have been damaged, for example, accessible *Polytrichum* and *Chorisodontium* moss banks and *Deschampsia* on the north-eastern and eastern sides of the island have been extensively damaged in some locations. A visit in February 2011 reported fur seals were present over the eastern side of the island [roughly drawing a line between the boat landing site (Lat. 60°39'05" S, Long. 045°36'12" W; Figure 2) and the island's summit (Lat. 60°39'05" S, Long. 045°36'12" W)]. Seals were present to the highest point of the island with about 30 seals on the summit. During a subsequent visit in January 2022, no fur seals were observed. Despite variable levels of fur seal trampling, both the Antarctic hair grass; *Deschampsia antarctica* and *Colobanthus quitensis* have thrived over recent years. The area covered by *Deschampsia*, as reported in February 2011, is more extensive than in the previous report (February 1999). The grass has now increased its abundance and distribution range in an area to the east of the island, extending west to the highest point on the island with good cover to the summit and all over the area around the summit cairn (Figure 3).

During a visit in February 1999 it was observed that the most luxuriant areas of grass on the northern and north-western slopes had not yet been affected and this observation was confirmed during a visit in February 2011. Notwithstanding some localised destruction, to date the primary values of the island, as noted above, have not been significantly compromised by either human or seal access to the island.

Resolution 3 (2008) recommended that the "Environmental Domains Analysis for the Antarctic Continent", be used as a dynamic model for the identification of Antarctic Specially Protected Areas within the systematic environmental-geographical framework referred to in Article 3(2) of Annex V of the Protocol (see also Morgan et al., 2007). ASPA 110 is not categorised within Morgan et al.; however, ASPA 110 is likely to be contained within Environment Domain G (Antarctic Peninsula off-shore islands geologic). The scarcity of Environment Domain G, relative to the other environmental domain areas, means that substantial efforts have been made to conserve the values found within this environment type elsewhere: other protected areas containing Domain G include ASPAs 109, 111, 112, 125, 126, 128, 145, 149, 150, and 152 and ASMAs 1 and 4.

Resolution 3 (2017) recommended that the Antarctic Conservation Biogeographic Regions (ACBRs) be used for the 'identification of areas that could be designated as Antarctic Specially Protected Areas within the systematic environmental-geographic framework referred to in Article 3(2) of Annex V to the Environmental Protocol. ASPA 110 sits within Antarctic Conservation Biogeographic Region (ACBR) 2 South Orkney Islands.

The two other ASPAs present within the South Orkney Islands (ASPA No. 109 Moe Island, and ASPA No. 111 Southern Powell Island and adjacent islands) were designated primarily to protect terrestrial vegetation and bird communities. ASPA No. 110 Lynch Island complements the local network of ASPAs by protecting a representative sample of the maritime Antarctic ecosystem including phanerogam-dominated terrestrial communities.



1. Description of values to be protected

Following a visit to the ASPA in January 2022, the values specified in the earlier designation were reviewed. Values within the Area are set out as follows:

- The Area contains luxuriant swards of Antarctic hair grass *Deschampsia antarctica* and the only other Antarctic flowering plant, Antarctic pearlwort (*Colobanthus quitensis*), is also abundant. It is also one of few sites where the grass *Deschampsia* is known to grow directly on *Polytrichum-Chorisodontium* moss banks.
- The cryptogamic vegetation is typical of the region; however, several species of moss found on the island (*Polytrichastrum alpinum* (= *Polytrichum alpinum*) and *Muelleriella crassifolia*) are unusually fertile for their southerly location. It is also possibly the only known location in Antarctica where *Polytrichastrum alpinum* develops sporophytes in profusion annually. Furthermore, *Polytrichum strictum* (= *Polytrichum alpestre*) occasionally produces male inflorescences in local abundance, which is a rare occurrence in this species in Antarctica and the rare moss *Plagiothecium ovalifolium* occurs in moist shaded rock crevices near the shore.
- The shallow loam-like soil associated with the grass swards contains a rich invertebrate fauna. The population density of the arthropod community associated with *Deschampsia* on Lynch Island appears unusually high, with some measurements suggesting it is one of the highest in the world. The site also shows unusual diversity for an Antarctic site. A rare enchytraeid worm was also found in moist moss in rock crevices on the northern side of the island. One arthropod species (*Globoppia loxolineata*) is near the northernmost limit of its known distribution, and specimens collected from Lynch Island exhibited unusual morphological characteristics compared to specimens collected elsewhere in the South Orkney-Antarctic Peninsula region.
- *Chromobacterium* bacteria, yeasts and fungi are found in higher densities than on Signy Island, thought to be a result of the lower acidity of the soils associated with *Deschampsia* and the more favourable microclimate at Lynch Island.
- The shallow gravelly loam-like soil beneath the dense swards of *Deschampsia* may represent one of the most advanced soil types in the Antarctic.

2. Aims and objectives

Management at Lynch Island aims to:

- avoid major changes to the structure and composition of the terrestrial vegetation;
- prevent unnecessary human disturbance to the Area;
- prevent or minimise the introduction to the Area of non-native plants, animals and microorganisms;
- allow scientific research in the Area provided it is for compelling reasons which cannot be served elsewhere and which will not jeopardise the natural ecological system in that Area;
- ensure that the flora and fauna are not adversely affected by excessive sampling within the Area;
- allow visits for management purposes in support of the aims of the management plan;
- minimise the possibility of introduction of pathogens which may cause disease in vertebrate populations within the Area;

3. Management activities

The following management activities shall be undertaken to protect the values of the Area

- Visits shall be made as necessary to assess whether the ASPA continues to serve the purposes for which it was designated and to ensure management and maintenance measures are adequate.
- The Management Plan shall be reviewed at least every five years and updated as required.
- Markers, signs or other structures erected within the Area for scientific or management purposes shall be secured and maintained in good condition and removed when no longer required.
- In accordance with the requirements of Annex III to the Protocol on Environmental Protection to the Antarctic Treaty, abandoned equipment or materials shall be removed to the maximum extent possible provided doing so does not adversely impact on the environment and the values of the Area.
- A copy of this Management Plan shall be made available at Signy Research Station (UK; 60°42'30" S, 045°36'30" W) and Orcadas Station (Argentina; 60°44'15" S, 044°44'20" W).
- Where appropriate, national Antarctic programmes are encouraged to liaise closely to ensure management activities are implemented. In particular, national Antarctic programmes are encouraged to consult with one another to prevent excessive sampling of biological material within the Area. Also, national Antarctic programmes are encouraged to consider joint implementation of guidelines intended to minimize the introduction and dispersal of non-native species within the Area.
- All scientific and management activities undertaken within the Area should be subject to an Environmental Impact Assessment, in accordance with the requirements of Annex I to the Protocol on Environmental Protection to the Antarctic Treaty.



4. Period of designation

Designated for an indefinite period.

5. Maps and images

Figure 1. Map of the location of Lynch Island in relation to the South Orkney Islands and the other protected areas in the region. Inset: the location of the South Orkney Islands in Antarctica. Map specifications: Projection: WGS84 Antarctic Polar Stereographic. Standard parallel: 71 °S. Central meridian 45 °W.

Figure 2. ASPA No. 110, Lynch Island, South Orkney Islands, topographic map. Projection: Lambert Conformal Conic. Standard parallels: 1st 60°40'00" W; 2nd 63°20'00" S. Central Meridian: 045°26'20" W. Latitude of Origin: 63°20'00" S. Spheroid: WGS84. Datum: Mean Sea Level. Horizontal accuracy of control points: ±1 m.

Figure 3. Normalised Difference Vegetation Index (NDVI), derived from satellite imagery, for ASPA No. 110 Lynch Island, South Orkney Islands, showing green vegetation cover using a colour scale of yellow → orange → red, with red indicating the highest NDVI values.

6. Description of the Area

6(i) Geographical co-ordinates, boundary markers and natural features

BOUNDARIES AND CO-ORDINATES

The Area encompasses all of Lynch Island but excludes all unnamed adjacent islands and islets. The Area encompasses all of the ice-free ground, permanent ice and semi-permanent ice found within Lynch Island, but excludes the marine environment extending greater than 10 m offshore from the low tide water line (Map 2). Boundary markers have not been installed because the coast itself is a clearly defined and visually obvious boundary.

GENERAL DESCRIPTION

Lynch Island (Latitude 60°39'10" S, Longitude 045°36'25" W; area) is a small island situated at the eastern end of Marshall Bay in the South Orkney Islands, about 200 m south of Coronation Island and 2.4 km north of Signy Island (Map 1). The 500 m x 300 m island has low cliffs of up to 20 m in height on the south, east and west sides, dissected by boulder-filled gullies. The northern side has a low cliff below a rock terrace at about 5-8 m altitude, above which moderate slopes rise to a broad plateau at about 40-50 m, with a maximum altitude of 57 m. A beach at the eastern end of the northern coast affords easy access to relatively gentle slopes leading to the central plateau area. The coastal cliffs generally make access to the upper island by other routes difficult, although access is feasible via one or two of the gullies on the eastern and northern sides. Small temporary melt-streams occur on the slopes in summer, but there are no permanent streams or pools, and only a few small late-lying snow patches occur on the southern side of the island. No meteorological data are available for Lynch Island, but conditions are broadly expected to be similar to those experienced at Signy Research Station. However, anecdotal observations suggest that significant microclimatic differences exist on Lynch Island, as the more profuse growth of plant communities would seem to attest.

The island is exposed to the south-west and to katabatic and föhn winds descending from Coronation Island to the north. However, in other respects the island is relatively sheltered from regional northerly, easterly and southerly winds by Coronation Island, Cape Hansen and Signy Island respectively. The föhn effect can briefly raise local air temperatures by as much as 10°C at Signy Island. Lynch Island has often been observed to receive sunshine when the surrounding region is shrouded in low cloud. The angle of solar incidence is also relatively high on the northern side of the island because of its general slope and aspect. The above factors may be important reasons for the abundance of the two flowering plants found on the island.

GEOLOGY

The bedrock of Lynch Island consists of quartzo-feldspathic and micaceous schists of the Scotia metamorphic complex, but is poorly exposed and equivalent rocks are much better displayed in the Cape Hansen area, to the east on Coronation Island.

PEDOLOGY

Three main soil types have been identified on Lynch Island:

- (i) An acidic (pH 3.8 – 4.5) moss peat, formed by the tall turf-forming mosses *Chorisodontium aciphyllum* and *Polytrichum strictum* (= *Polytrichum alpestre*), occurs mainly at the north-eastern end of the island. This peat reaches a depth of about 50 cm and is similar to peat on Signy Island where it reaches a depth of 2 m. Where the peat depth exceeds about 30 cm there is permafrost. In a few places where the substratum is moist, shallow peat of 10-15 cm depth (pH 4.8 - 5.5) has accumulated beneath the carpet-forming mosses *Warnstorfia laculosa* (= *Calliergidium austro-stramineum*) and *Sanionia uncinata* (= *Drepanocladus uncinatus*).
- (ii) A shallow, gravelly loam-like soil resembling tundra brown soil occurs beneath dense swards of the grass *Deschampsia antarctica*. It is seldom more than about 30 cm in depth (pH 5.0 – 5.8) and probably represents one of the most advanced soil types in the Antarctic.
- (iii) A glacial till with material ranging from fine clay (pH 5.2 – 6.0) and sand to gravel and larger stones. This covers the summit plateau and occurs in rock depressions throughout the island, as well as on parts of the rock terrace. On the plateau cryoturbation has in several places sorted the material into patterned features with small stone circles and polygons on level ground and stone stripes on sloping ground. At the north-eastern end of the island, the deposition of limpet shells (*Nacella concinna*) by gulls (*Larus dominicanus*) has resulted in a more calcareous mineral soil in rock



TERRESTRIAL FLORA

Cryptogamic and phanerogamic vegetation typical of the maritime Antarctic is found over much of the island (Figure 3). Use of satellite remote sensing techniques (Normalised Difference Vegetation Index) showed the area of green vegetation within the ASPA to be 35,000 m² (25% of the ASPA area). The most significant aspect of the vegetation is the abundance and reproductive success of the two native Antarctic flowering plants, the Antarctic hair grass (*Deschampsia antarctica*) and Antarctic pearlwort (*Colobanthus quitensis*), found especially on the northern slopes (Map 3). Both species flower in profusion and seed viability appears to be much greater than on Signy Island. Lynch Island possesses the largest stands of *Deschampsia* and the greatest abundance of *Colobanthus* known in the South Orkney Islands and one of the most extensive anywhere in the Antarctica Treaty area. On the rock terrace and moist slope rising above the northern coast, the grass forms extensive swards of up to 15 × 50 m. These swards range from continuous stands of relatively luxuriant plants on the moister sites and ledges to small, yellowish, more isolated plants on the drier, stonier and more exposed terrain. *Colobanthus* is generally associated with the grass, but here the plants do not coalesce to form closed patches. This is one of very few sites where *Deschampsia* is known to grow directly on *Polytrichum-Chorisodontium* moss banks. Elsewhere on the island, the grass and, to a lesser extent, the pearlwort are frequent associates in other communities, especially stands of denser fellfield vegetation where there is quite high cover afforded by various mosses and lichens (particularly towards the western end of the northern terrace).

Shallow but occasionally extensive (about 50 m²) banks of *Chorisodontium aciphyllum* and *Polytrichum strictum* are frequent at the north-eastern end of the island and, to a lesser extent, on the southern side. These are typical of the moss banks which occur on Signy Island and elsewhere in the northern maritime Antarctic, with several fruticose and crustose lichens growing epiphytically on the moss surface. In small moist depressions, there are carpets of *Warnstorfia laculosa* and *Sanionia uncinata*, with some *Warnstorfia sarmentosa* (= *Calliergon sarmentosum*) and *Cephaloziella varians* (= *C. exiliflora*). On wet soil and rock ledges, *Brachythecium austro-salebrosum* is common. On the drier, more windswept, stonier soils and rock surfaces – notably in the plateau area – a typical open fellfield community of many bryophyte and lichen taxa form a complex mosaic. The dominant species in this locality are the lichens *Usnea antarctica* and *U. aurantiaco-atra* (= *U. fasciata*) and the moss *Andreaea depressinervis*; *Sphaerophorus globosus* and other species of *Alectoria*, *Andreaea*, *Cladonia*, and *Stereocaulon* are also common, while *Himantormia lugubris* and *Umbilicaria antarctica* are infrequent. Crustose lichens are abundant on all rock surfaces. The mosses and macrolichens in this area are loosely attached on thin soils and are easily damaged. Large thalli of *Usnea* spp. and *Umbilicaria antarctica* are found on moist sheltered boulders and rock faces, especially on the southern side of the island.

Communities of crustose lichen occur in the cliffs above the high water mark, especially where the rock is influenced by breeding or roosting birds. The distribution of several species forms distinctive zones in relation to inundation by sea spray and exposure to wind. The best developed communities of brightly coloured ornithocrophilous taxa occur at the western end of the island where *Caloplaca* spp., *Haematomma erythromma*, *Mastodia tessellata*, *Physcia caesia*, *Xanthoria candelaria*, *X. elegans*, and species of *Buellia* and *Verrucaria* are frequent. The uncommon halophilous moss *Muelleriella crassifolia* also occurs within the spray zone around the island.

The only rare moss recorded on Lynch Island is *Plagiothecium ovalifolium*, found in moist, shaded rock crevices near the shore. However, the island is possibly the only site known in the Maritime Antarctic where the moss *Polytrichastrum alpinum* develops sporophytes in profusion each year; this occurs among *Deschampsia*, *Colobanthus* and cryptogams on the northern side of the island; elsewhere in the Antarctic sporophytes are in some years very rare. Also, *Polytrichum strictum* produces male inflorescences in local abundance, a rare phenomenon in this species in the Antarctic. While the thalloid liverwort *Marchantia berteroana* is locally common on Signy Island, Lynch Island is one of very few other localities where it is known in the South Orkney Islands. Several cryptogamic species of very restricted distribution in the Antarctic, but which are locally common on Signy Island and the mainland of Coronation Island only a few hundred metres away, have not been observed at Lynch Island.

TERRESTRIAL INVERTEBRATES

The microinvertebrate fauna associated with the rich *Deschampsia* swards described thus far comprises 13 taxa: three springtails (*Cryptopygus antarcticus*, *Friezea woyciechowskii* and *Isotoma (Folsomotoma) octooculata* (= *Parisotoma octooculata*), one mesostigmatid mite (*Gamasellus racovitza*), two cryptostigmatid mites (*Alaskozetes antarcticus* and *Globoppia loxolineata*), and seven prostigmatid mites (*Apotriphyteus* sp., *Ereynetes macquariensis*, *Nanorchestes berryi*, *Stereotydeus villosus*, and three species of *Eupodes*). The number of taxa identified is likely to increase with greater sampling. The community is dominated by the Collembolla, especially *Cryptopygus antarcticus* (84% of all arthropods extracted), with relatively large numbers of *I. octooculata*; the principal mite was an undetermined species of *Eupodes*. *Globoppia loxolineata* is near the northernmost limit of its known distribution. In general, the population density of the arthropod community of grass stands on Lynch Island appears unusually high, with some measurements suggesting it is one of the highest in the world. It also shows considerable diversity for an Antarctic site, although this observation was based on a small number of sample replicates and further sampling would be required to establish densities with greater reliability: this is difficult to achieve on Lynch Island given the very limited extent of communities available for sampling.



Lynch Island was the first site in the Antarctic where a terrestrial enchytraeid was found (in soil beneath a moss *Hennediella antarctica* on a rock ledge above the northern shore); only in a few other sites in the South Orkney Islands have these worms been found – although few samples have been gathered and the species has yet to be identified. Of the tardigrade fauna, most of the 16 individuals isolated from a sample of *Brachythecium* were *Hypsibius alpinus* and *H. pinguis* with some *H. dujardini*, while of 27 isolated from a *Prasiola crispa* sample, almost all were the latter species with a few that were other species of *Hypsibius*.

MICROORGANISMS

The mineral and organic soils of Lynch Island have a slightly higher pH than corresponding soils on nearby Signy Island. This higher base and nutrient status, together with the more favourable microclimate, is reflected in larger numbers of bacteria (including *Chromobacterium*), yeasts and fungi than occur in comparable soils on Signy Island. Bacterial numbers in the *Polytrichum* peat on Lynch Island are about eight times, and in the *Warnstorfia* peat about six times, greater than in corresponding Signy Island peats; yeasts and fungi are similarly much more abundant. Soil associated with the two flowering plants yielded several nematophagous fungi: in *Deschampsia* soil *Acrostalagmus goniodes*, *Cephalosporium balanoides* and *Dactylaria gracilis*; in *Colobanthus* soil, *Cephalosporium balanoides*, *Dactylaria gracilis*, *Dactylella stenobrocha* and *Harposporium anguillulae* were found. The basidiomycete fungi *Galerina antarctica* and *G. longinqua* occur on moist moss.

VERTEBRATES

The island has no penguin colonies or substantial breeding colonies of other birds. Groups of chinstrap (*Pygoscelis antarctica*), Adélie (*P. adeliae*) and gentoo (*P. papua*) penguins and, sometimes, blue-eyed cormorants (*Phalacrocorax atriceps*) often congregate at the north-eastern and the western ends of the island. Several pairs of brown skuas (*Catharacta lonnbergii*) and at least two pairs of kelp gulls (*Larus dominicanus*) were observed in the early 1980s to nest at the north-eastern corner. A small colony of Antarctic terns (*Sterna vittata*) may also occur in this vicinity, although in February 1994 breeding was not observed. Cape petrels (*Daption capense*) and snow petrels (*Pagodroma nivea*) breed on the higher cliffs at the eastern end and along the north-western coast of the island. A few pairs of snow petrels and Wilson's storm petrels (*Oceanites oceanicus*) nest on ledges and beneath boulders on the south side of the island.

Weddell seals (*Leptonychotes weddellii*), crabeater seals (*Lobodon carcinophagus*), occasional leopard seals (*Hydrurga leptonyx*), and small groups of southern elephant seals (*Mirounga leonina*) are regularly seen on the coast and on ice floes in the vicinity; none have been known to breed on Lynch Island. Since the early 1980s increasing numbers of Antarctic fur seals (*Arctocephalus gazella*), virtually all being immature non-breeding males, have been observed on Lynch Island, some gaining access up the more gentle north-eastern slopes to vegetated areas, where they have caused local, but severe, damage to *Polytrichum-Chorisodontium* moss banks and other communities.

Seal access to the island is principally from a beach on the northeast coast. Once seals have gained access, there are no further substantial geographical impediments to their more extensive travel over the island. Groups of seals have been observed near the summit. Destruction of swards of *Deschampsia* was first reported in 1988. During earlier inspections of the island, it was observed that the most luxuriant areas of *Deschampsia* and *Colobanthus* on the northern and north-western slopes had not yet been affected. Accessible areas of vegetation in the eastern and north-eastern sides of the island, particularly *Polytrichum* and *Chorisodontium* moss banks, had been severely damaged by Antarctic fur seals. In some eastern and north-eastern areas that have been heavily impacted by fur seals, *Deschampsia* and *Colobanthus* have either been damaged or have died, but at less impacted locations at higher altitudes, these plants continue to grow and may be increasing their abundance and extending their distribution range on the island (see Map 3). During the most recent inspection, no fur seals were observed on the island.

6(ii) Access to the Area

- Where possible, access shall be by small boat. Landings from the sea should be at the beach on the eastern end of the northern coast of the island (Lat. 60°39'05" S, Long. 045°36'12" W; Map 2), unless specifically authorised by Permit to land elsewhere, or when landing at this location is impractical because of adverse conditions.
- Under exceptional circumstances, necessary for purposes consistent with the objectives of the Management Plan, helicopters may be permitted to land within the Area.
- Landing of helicopters within the Area shall be at the designated location on the rock platform (8 m) on the north-western end of the island (Lat. 60°39'04.5" S, Long. 045°36'12" W; Map 2).
- Within the Area the operation of aircraft should be carried out, as a minimum requirement, in compliance with the 'Guidelines for the Operation of Aircraft near Concentrations of Birds' contained in Resolution 2 (2004). When conditions require aircraft to fly at lower elevations than recommended in the guidelines, aircraft should maintain the maximum elevation possible and minimise the time taken to transit.
- Use of helicopter smoke grenades is prohibited within the Area unless absolutely necessary for safety. If used, all smoke grenades should be retrieved.

6(iii) Location of structures within and adjacent to the Area

There are no structures present in the Area apart from several cairns marking sites used for topographical survey. The island's summit cairn is located at Lat. 60°39'05" S, Long. 045°36'12" W. A sign notifying the protected status of Lynch Island was erected on a prominent rock outcrop above the recommended landing beach in February 1994, but this was destroyed by strong winds.

Signy Research Station (UK) is 6.4 km south at Factory Cove, Borge Bay, on Signy Island.



6(iv) Location of other protected areas in the vicinity

The nearest protected areas to Lynch Island are Moe Island (ASPA No. 109), which is about 10 km SSW, and Southern Powell Island and adjacent islands (ASPA No. 111), which is about 35 km to the east (Map 1).

6(v) Special zones within the Area

None.

7. Permit conditions

7(i) General permit conditions

Entry into the Area is prohibited except in accordance with a Permit issued by an appropriate national authority as designated under Article 7 of Annex V of the Protocol on Environmental Protection to the Antarctic Treaty.

Conditions for issuing a Permit to enter the Area are that:

- it is issued for a compelling scientific purpose which cannot be served elsewhere; or
- it is issued for essential management purposes such as inspection, maintenance or review;
- the actions permitted will not jeopardise the natural ecological system in the Area;
- any management activities are in support of the objectives of this Management Plan;
- the actions permitted are in accordance with this Management Plan;
- the Permit, or an authorised copy, must be carried within the Area;
- permits shall be issued for a stated period;
- a report or reports are supplied to the authority or authorities named in the Permit;
- the appropriate authority should be notified of any activities/measures undertaken that were not included in the authorised Permit.

7(ii) Access to, and movement within or over, the Area

- Land vehicles are prohibited within the Area
- Movement within the Area shall be on foot.
- Pilots, helicopter or boat crew, or other people on helicopters or boats, are prohibited from moving on foot beyond the immediate vicinity of their landing site unless specifically authorised by Permit.
- Pedestrian traffic should be kept to the minimum consistent with the objectives of any permitted activities and every reasonable effort should be made to minimise trampling effects, i.e. all movement should be undertaken carefully so as to minimise disturbance to the soil and vegetated surfaces, walking on rocky terrain if practical.
- Overflight of bird colonies within the Area by Remotely Piloted Aircraft Systems (RPAS) shall not be permitted unless for compelling scientific or operational purposes, and in accordance with a permit issued by an appropriate national authority. Furthermore, operation of RPAS within or over the Area shall be in accordance with the 'Environmental guidelines for operation of Remotely Piloted Aircraft Systems (RPAS) in Antarctica' (Resolution 4 (2018)) (available at: https://documents.ats.aq/recatt/att645_e.pdf).

7(iii) Activities which may be conducted in the Area

- Compelling scientific research which cannot be undertaken elsewhere and which will not jeopardize the ecosystem of the Area
- Essential management activities, including monitoring

7(iv) Installation, modification or removal of structures

No new structures are to be erected within the Area, or scientific equipment installed, except for compelling scientific or management reasons and for a pre-established period, as specified in a permit. Installation (including site selection), maintenance, modification or removal of structures and equipment shall be undertaken in a manner that minimises disturbance to the values of the Area. All structures or scientific equipment installed in the Area shall be clearly identified by country, name of the principal investigator and year of installation. All such items should be free of organisms, propagules (e.g., seeds, eggs) and non-sterile soil (see Section 7(vi)), and be made of materials that can withstand the environmental conditions and pose minimal risk of contamination of the Area. Removal of specific structures or equipment for which the Permit has expired shall be a condition of the Permit. Permanent structures or installations are prohibited.



7(v) Location of field camps

Camping should be avoided within the Area. However, when absolutely necessary for purposes specified in the Permit, camping is allowed at the designated site at the north-western end of the island (Lat. 60°39'04" S, Long. 045°36'37" W; Map 2).

7(vi) Restrictions on materials and organisms which may be brought into the Area

No living animals, plant material or microorganisms shall be deliberately introduced into the Area. To ensure that the floristic and ecological values of the Area are maintained, special precautions shall be taken against accidentally introducing microbes, invertebrates or plants from other Antarctic sites, including stations, or from regions outside Antarctica. All sampling equipment or markers brought into the Area shall be cleaned or sterilized. To the maximum extent practicable, footwear and other equipment used or brought into the Area (including bags or backpacks) shall be thoroughly cleaned before entering the Area. Further guidance can be found in the *CEP non-native species manual* (Resolution 4 (2016)) and the *SCAR Environmental code of conduct for terrestrial scientific field research in Antarctica* (Resolution 5 (2018)).

No herbicides or pesticides shall be brought into the Area. Any other chemicals, including radio-nuclides or stable isotopes, which may be introduced for scientific or management purposes specified in the Permit, shall be removed from the Area at or before the conclusion of the activity for which the Permit was granted. Release of radio-nuclides or stable isotopes directly into the environment in a way that renders them unrecoverable should be avoided. Fuel or other chemicals shall not be stored in the Area unless specifically authorised by Permit condition. They shall be stored and handled in a way that minimises the risk of their accidental introduction into the environment. Materials introduced into the Area shall be for a stated period only and shall be removed by the end of that stated period. If release occurs which is likely to compromise the values of the Area, removal is encouraged only where the impact of removal is not likely to be greater than that of leaving the material in situ. The appropriate authority should be notified of anything released and not removed that was not included in the authorised Permit.

7(vii) Taking, or harmful interference with, native flora or fauna

Taking or harmful interference with native flora or fauna is prohibited, except by Permit issued in accordance with Annex II to the Protocol on Environmental Protection to the Antarctic Treaty. Where taking or harmful interference with animals is involved, the *SCAR Code of Conduct for the Use of Animals for Scientific Purposes in Antarctica* (Resolution 4 (2019)) should be used as a minimum standard.

7(viii) The collection or removal of materials not brought into the Area by the Permit holder

Collection or removal of anything not brought into the Area by the permit holder shall only be in accordance with a Permit and should be limited to the minimum necessary to meet scientific or management needs.

Permits shall not be granted if there is a reasonable concern that the sampling proposed would take, remove or damage such quantities of soil, native flora or fauna that their distribution or abundance within the Area would be significantly affected.

Other material of human origin likely to compromise the values of the Area which was not brought into the Area by the permit holder or otherwise authorised, may be removed from the Area unless the environmental impact of the removal is likely to be greater than leaving the material in situ; if this is the case the appropriate Authority must be notified and approval obtained.

7(ix) Disposal of waste

As a minimum standard, all waste shall be disposed of in accordance with Annex III to the Protocol on Environmental Protection to the Antarctic Treaty. In addition, all wastes shall be removed from the Area. Liquid human wastes may be disposed of into the sea. Solid human waste should not be disposed of to the sea, but shall be removed from the Area. No solid or liquid human waste shall be disposed of inland.

7(ix) Measures that may be necessary to continue to meet the aims of the Management Plan

- Permits may be granted to enter the Area to carry out scientific research, monitoring and site inspection activities, which may involve the collection of a small number of samples for analysis, to erect or maintain signboards, or to carry out protective measures.
- Any long-term monitoring sites shall be appropriately marked and the markers or signs maintained.
- Scientific activities shall be performed in accordance with the *SCAR Environmental Code of Conduct for Terrestrial Scientific Field Research in Antarctica* (Resolution 5 (2018)). Geological research shall be undertaken in accordance with the *SCAR Environmental Code of Conduct for Geosciences Field Research Activities in Antarctica* (Resolution 1 (2021)).

7(xi) Requirements for reports

The principal permit holder for each visit to the Area shall submit a report to the appropriate national authority as soon as practicable, and no later than six months after the visit has been completed. Such reports should include, as appropriate, the information identified in the visit report form contained in the Guide to the Preparation of Management Plans for Antarctic Specially Protected Areas. If appropriate, the national authority should also forward a copy of the visit report to the Party that proposed the Management Plan, to assist in managing the Area and reviewing the Management Plan. Wherever possible, Parties should deposit the original or copies of the original visit reports, in a publicly accessible archive to maintain a record of usage, for the purpose of any review of the Management Plan and in organising the scientific use of the Area.



8. Supporting documentation

Convey, P. 1994. Modelling reproductive effort in sub- and maritime Antarctic mosses. *Oecologia* **100**: 45-53.

Block, W. and Christensen, B. 1985. Terrestrial Enchytraeidae from South Georgia and the Maritime Antarctic. *British Antarctic Survey Bulletin* **69**: 65-70.

Bonner, W.N. and Smith, R.I.L. (Eds) 1985. *Conservation areas in the Antarctic*. SCAR, Cambridge: 73-84.

Bonner, W.N. 1994. Active management of protected areas. In Smith, R.I.L., Walton, D.W.H. and Dingwall, P.R. (Eds) *Developing the Antarctic Protected Area system. Conservation of the Southern Polar Region I*. IUCN, Gland and Cambridge: 73-84.

Booth, R.G., Edwards, M. and Usher, M.B. 1985. Mites of the genus *Eupodes* (Acari, Prostigmata) from maritime Antarctica: a biometrical and taxonomic study. *Journal of the Zoological Society of London (A)* **207**: 381-406. (samples of *Eupodes* analysed)

Burn, R. and Usher, M.B. 1986. A morphometric study of the mite, *Oppia loxolineata*, in the Maritime Antarctic. *British Antarctic Survey Bulletin* **73**: 47-50.

Chalmers, M.O. 1994. Lynch Island fur seal enclosure report 01/01/94. Unpublished British Antarctic Survey report BAS Ref AD6/2H/1993/NT2.

Greene, D.M and Holtom, A. 1971. Studies in *Colobanthus quitensis* (Kunth) Bartl. and *Deschampsia antarctica* Desv.: III. Distribution, habitats and performance in the Antarctic botanical zone. *British Antarctic Survey Bulletin* **26**: 1-29.

Hodgson, D.A. and Johnston, N.M. 1997. Inferring seal populations from lake sediments. *Nature* **387**(1 May).

Hodgson, D.A., Johnston, N.M., Caulkett, A.P., and Jones, V.J. 1998. Palaeolimnology of Antarctic fur seal *Arctocephalus gazella* populations and implications for Antarctic management. *Biological Conservation* **83**(2): 145-54.

Hooker, T.N. 1974. Botanical excursion to Lynch Island, 13/03/74. Unpublished British Antarctic Survey report BAS Ref AD6/2H/1973-74/N12.

Hughes, K. A., Ireland, L., Convey, P., Fleming, A. H. 2016. Assessing the effectiveness of specially protected areas for conservation of Antarctica's botanical diversity. *Conservation Biology*, **30**: 113-120.

Jennings, P.G. 1976. Tardigrada from the Antarctic Peninsula and Scotia Ridge region. *British Antarctic Survey Bulletin* **44**: 77-95.

SCAR (Scientific Committee on Antarctic Research). 2009. Environmental code of conduct for terrestrial scientific field research in Antarctica. ATCM XXXII IP4.

Shears, J.R. and Richard, K.J. 1994. Marking and inspection survey of Specially Protected Areas in the South Orkney Islands, Antarctica 07/01/94 – 17/02/94. Unpublished British Antarctic Survey report BAS Ref AD6/2H/1993/NT5.

Smith, R.I. Lewis 1972. Vegetation of the South Orkney Islands. *BAS Scientific Report* **68**, British Antarctic Survey, Cambridge.

Smith, R.I. Lewis 1990. Signy Island as a paradigm of environmental change in Antarctic terrestrial ecosystems. In K.R. Kerry and G. Hempel. *Antarctic Ecosystems: ecological change and conservation*. Springer-Verlag, Berlin: 32-50.

Smith, R.I. Lewis 1994. Introduction to the Antarctic Protected Area System. In Smith, R.I.L., Walton, D.W.H. and Dingwall, P.R. (Eds) *Developing the Antarctic Protected Area system. Conservation of the Southern Polar Region I*. IUCN, Gland and Cambridge: 14-26.

Smith, R.I. Lewis 1997. Impact of an increasing fur seal population on Antarctic plant communities: resilience and recovery. In Battaglia, B. Valencia, J. and Walton, D.W.H. *Antarctic communities: species, structure and survival*. Cambridge University Press, Cambridge: 432-36.

Star, J. and Block, W. 1998. Distribution and biogeography of oribatid mites (Acari: Oribatida) in Antarctica, the sub-Antarctic and nearby land areas. *Journal of Natural History* **32**: 861-94.

Usher, M.B. and Edwards, M. 1984. The terrestrial arthropods of the grass sward of Lynch Island, a specially protected area in Antarctica. *Oecologia* **63**: 143-44.

Usher, M.B. and Edwards, M. 1986. A biometrical study of the family Tydeidae (Acari, Prostigmata) in the Maritime Antarctic, with descriptions of three new taxa. *Journal of the Zoological Society of London (A)* **209**: 355-83.

Wynn-Williams, D.D. 1982. The microflora of Lynch Island, a sheltered maritime Antarctic site. *Comité National Française Recherche en Antarctiques* **51**: 538.



Figure 1. Map showing the location of Lynch Island in relation to the South Orkney Islands and the other protected areas in the region. Inset: the location of the South Orkney Islands in Antarctica.




Figure 2. ASPA No. 110, Lynch Island, South Orkney Islands, topographic map.






Figure 3. Normalised Difference Vegetation Index (NDVI), derived from satellite imagery, for ASPA No. 110 Lynch Island, South Orkney Islands, showing green vegetation cover using a colour scale of yellow → orange → red, with red indicating the highest NDVI values

